

CHEMICAL THREAT TO CHILDREN

By

Murray Thompson

(BappSci Environmental Health 1998, University of Western Sydney)



HERBICIDES COMMONLY USED ON SCHOOL GROUNDS

In point form:

- Glyphosate has been described as “extremely persistent” (NCAP 2000:2, citing US EPA 1993-2), even though it has been promoted as an environmentally friendly herbicide that rapidly breaks down. However, tests show that glyphosate may persist for 3 years (NCAP 2000:2, citing Torstensson et al. 1989), while its metabolite, AMPA, may persist even longer (Ewing 1999:11, citing World Health Organization 1994). Glyphosate has been shown to cause genetic mutations in human, animal and plant cell tests (NCAP 2000:2, citing Vigfusson 1980; Kale et al. 1995; Rank et al. 1993).
- Oryzalin (active ingredient of Surflan) is persistent and carcinogenic (NCAP 2000:2, citing US EPA 1994).
- Dichlobenil (active ingredient of Casoron) is persistent (NCAP 2000:2, citing Williams & Eagle 1979), causes cancer in animals and is classified by EPA as a possible human carcinogen (NCAP 2000:2, citing US EPA 1999-1).
- Glyphosate has been associated with a number of health and ecological problems (Cox 1993:4). The surfactants added to glyphosate are acutely toxic to humans and aquatic animals, can cause damage to the digestive system and lungs, and may be the cause of death in exposed humans (Cox 1993:4, citing Sawada, et al. 1988; Tominack 1991; Talbot 1991).
- Glyphosate ingestion has affected the pituitary gland and kidneys, and caused abnormal bone development and decreased birth weights in laboratory animals (Cox 1993:4, citing US EPA 1986).
- And, though the US EPA classified glyphosate as Group E, Evidence of Noncarcinogenicity in Humans, Monsanto’s own data submitted to EPA in support of this classification showed otherwise, note:
 - * increase in pancreatic tumors at 2 doses in female rats;
 - * significant increase in liver tumors with increasing dose;


* significant increase in thyroid tumors with increased dose in female rats (Cox 1993:4, citing Dykstra & Ghali 1991).

MISDIAGNOSIS AND SUSCEPTIBILITY TO CHEMICAL HARM

Two significant problems exist with regard to herbicide or pesticide poisoning:

1. Pesticide poisoning is extremely problematic because mild to moderate pesticide poisoning symptoms can be easily misdiagnosed as stomach-flu, bronchitis or asthma (Reeves, Schafer, Hallward & Katten 1999:17). My concern here is that some instances of flu and other outbreaks at local schools can be directly attributed to the spraying of chemicals, and are not simply issues of the regular childhood contagions we expect to see every year. Blood tests can confirm the presence or absence of toxins. They are a valuable medical avenue through which a situation can be clarified.

Even severe pesticide poisoning in infants has been misdiagnosed as aneurysm, head trauma, diabetic acidosis, severe bacterial gastroenteritis, pneumonia and whooping cough (Reeves, Schafer, Hallward & Katten 1999:17, citing Solomon & Mott 1998; Zweinerd & Ginsburg 1988).



<http://www.referralware.com/home.jsp/1398481110>

2. Children are much more susceptible to poisoning than adults for a number of reasons:

- * They inhale a greater volume of air in relation to their body weight than adults, and so receive a larger dose (NCAP 2000:12-14, citing NRC 1993).
- * They have a greater skin surface area proportional to body mass. This means that equivalent exposures in an adult and child results in a greater dose for the child (NCAP 2000:12-14, citing NRC 1993).
- * They play nearest the sites of application (NCAP 2000:13).
- * They display hand-to-mouth behavior (NCAP 2000:12,14).
- * Children's nervous systems are more susceptible to the impact of nerve poisons (NCAP 2000:12,14:4, citing NRC 1993; Watanabe 1990).
- * Children cannot effectively detoxify certain chemicals (compared to adults) and so are much more vulnerable to poisoning (NCAP 2000:12,14, citing NRC 1993).
- * The dividing cells in children's bodies are more susceptible to the impact of cancer-causing chemicals (NCAP 2000:12,14, citing NRC 1993).
- * Their immune systems, being underdeveloped, are more prone to damage from exposure to foreign compounds (NCAP 2000:12,14, citing Repetto & Baliga 1996; NRC 1993).

And also, in point form, regarding commonly used pesticides such as organophosphates (OPs):

- Organophosphate (OP) insecticides can be inhaled, whereupon they are rapidly absorbed (NCAP 2000:11).
- OPs are neurotoxicants, and are known to inhibit the enzyme acetylcholinesterase (Environmental Protection Agency 1995:26938).
- Human and experimental animal evidence exists "indicating that there may be residual, if not permanent, effects of exposure" (Environmental Protection Agency 1995, citing: Steenland et al. 1994; Tandon et al. 1994; Stephens et al. 1995).
- OP pesticides have been implicated in bone abnormalities in fish (Ewing 1999:35).
- The OP poisoning of children (including ingestion *and* inhalation of the chemicals) has resulted in the generation of numerous symptoms and events including slowed

heart rate, pulmonary edema (where some of the children required ventilators for several hours), death from parathion poisoning, stupor, coma, hypotonicity¹ and muscle weakness, and some muscarinic² symptoms (Pesticide Action Network 1999:2, citing Lifshitz et al. 1999:102-103).

- The OP poisoning of farm families (not themselves occupationally exposed, but merely living with farmers who used OPs) in agricultural communities in rural El Salvador presented symptoms such as limb cramps, chest pressure, changes in defecation, feeling dazed and increased lacrimation – all acute health effects produced in individuals not involved in field work themselves (Pesticide Action Network 1999:4, citing Azaroff & Neas 1999:158-164)).



- Home fumigation with OPs has produced acute and reversible parkinsonism (symptoms of Parkinson's Disease, or a syndrome resembling Parkinson's) (Pesticide Action Network 1999:4, citing Bhatt et al. 1999:1467-1471).
- Animal study evidence suggests that OP pesticide exposure can lead to neurodevelopmental effects such as reduced balance, increasing righting reflex time, and diminished cliff avoidance (Pesticide Action Network 2000:2, citing Eskenazi et al. 1999:409-416).
- Occupational exposure to OPs in China resulted in increases to rates of sperm aneuploidy (Pesticide Action Network 2000:4, citing Padungtod et al. 1999:230-238).
- **OP pesticides** (eg oxydemeton-methyl, methyl parathion and methamidophos) are **"responsible for most of the occupational deaths and poisonings in the U.S. and throughout the world"** (Reeves, et al. 1999:18, citing Blondell & Dobozy 1997; Keifer & Mahurin 1997; Moses et al. 1993, Savage et al. 1988).

¹ *Hypotonia*: Loss of muscular tonicity, relaxation of the arteries (Stedman's Medical Dictionary 1976:683).

² *Muscarinic*: Cardiac inhibition, vasodilation, salivation, lacrimation, bronchoconstriction, gastrointestinal stimulation (Stedman's Medical Dictionary 1976:893).

The above information demonstrates clearly that the Roundup or glyphosate herbicide is extremely toxic. Pesticides are also chemicals that need to be restricted far more than they generally are. This well researched and catalogued American data alone warrants a careful review of all pesticides and herbicides used in Australian schools.

REFERENCES

Azaroff, L.S. & Neas, L.M. 1999, 'Acute health effects associated with nonoccupational pesticide exposure in rural El Salvador', *Environmental Research (Section A)*, 80(2), February.

Bhatt, M.H. et al. 1999, 'Acute and reversible Parkinsonism due to organophosphate pesticide intoxication: five cases', *Neurology*, 52(7), April 22.

Blondell, J. & Dobozy, V.A. 1997, 'Review of chlorpyrifos poisoning data', January 1997 memorandum, Office of Prevention, Pesticides and Toxic Substances, US EPA, Washington, D.C., USA.

Cox, C. 1993, 'Biotechnology and Agricultural Pesticide Use: An Interaction Between Genes and Poisons', *Journal of Pesticide Reform*, Vol. 13, No. 3 (Fall), NCAP, PO Box 1393, Eugene, OR 97440, USA.

Cox, C. 1998, 'Jimmy and Jane's day: a precautionary tale', in *Journal of Pesticide Reform*, Summer 1998 – Vol. 18, No. 2, Northwest Coalition for Alternatives to Pesticides, PO Box 1393 Eugene, OR 97440-1393 USA (www.pesticide.org).

Dykstra, W. & Ghali, G.Z. 1991, 'Second peer review of glyphosate. Memo to R. Taylor and L. Rossi', US EPA Office of Pesticides and Toxic Substances, Health Effects Division, Washington, D.C.

Environmental Protection Agency 1995, 'Proposed Guidelines for Neurotoxicity Risk Assessment', *Federal Register* (Part II), Vol. 63, No.93.

Eskenazi, B. et al. 1999, 'Exposures of children to organophosphate pesticides and their potential adverse health effects', *Environmental Health Perspectives*, 107 (3), June.

Ewing, R.D. 1999, *Diminishing Returns: Salmon Decline and Pesticides*, Oregon Pesticide Education Network.

Keifer, M.C. & Mahurin, R.K. 1997, 'Chronic neurologic effects of pesticide overexposure', *Occup. Med.: State of the Art Reviews*, 12:291-304.



Kale, et al. 1995, 'Mutagenicity testing of nine herbicides and pesticides currently used in agriculture', *Environ. Mol. Mutagen.* 25:148-153.

Lifshitz, M. et al. 1999, 'Carbamate and organophosphate poisoning in young children', *Pediatric Emergency Care* 15(2), April.

Moses, M., Johnson, E.S., Anger, W.K., Burse, V.W., Horstman, S.F., Jackson, R.J., Lewis, R.G., Maddy, K.T., McConnell, R., Meggs, W.J., and Zahm, S.H. 1993, 'Environmental equity and pesticide exposure', *Toxicol. Indust. Health*, 9:913-959.

NCAP (Northwest Coalition for Alternatives to Pesticides) 1998, *Worst Kept Secrets: Toxic Inert Ingredients in Pesticides*, Northwest Coalition for Alternatives to Pesticides, PO Box 1393 Eugene, OR 97440-1393 USA (www.pesticide.org).

NCAP (Northwest Coalition for Alternatives to Pesticides) 2000, *Unthinkable Risk: How Children Are Exposed and Harmed When Pesticides Are Used at School*, Northwest Coalition for Alternatives to Pesticides, PO Box 1393 Eugene, OR 97440-1393 USA (www.pesticide.org).

NCAP (Northwest Coalition for Alternatives to Pesticides) 2000a, *Unintended Casualties: Five Children Whose Lives Were Profoundly Affected by Pesticide Exposures at School*, Northwest Coalition for Alternatives to Pesticides, PO Box 1393 Eugene, OR 97440-1393 USA (www.pesticide.org).

NRC 1993, *Pesticides in the diets of infants and children*, National Research Council, National Academy Press, Washington, DC.

Padungtod, C. et al. 1999, 'Sperm aneuploidy among Chinese pesticide factory workers: scoring by the FISH method', *American Journal of Industrial Medicine*, 36:2, August.

Pesticide Action Network (North America) 1999, *Pesticide Research Updates*, October, Number 1.

Pesticide Action Network (North America) 2000, *Pesticide Research Updates*, February, Number 2.

Rank, et al. 1993, 'Genotoxicity testing of the herbicide Roundup and its active ingredient glyphosate isopropylamine using the mouse bone marrow micronucleus test', *Salmonella* mutagenicity test and *Allium* anaphase-telophase test, *Mutation Research* 300:29-36.

Reeves, M., Schafer, K., Hallward, K. & Katten, A. 1999, *Fields of Poison: California Farmworkers and Pesticides*, Pesticide Action Network North America Regional Center,

California Rural Legal Assistance Foundation, United Farm Workers of America and Californians for Pesticide Reform (Publishers).

Repetto & Baliga 1996, *Pesticides and the Immune System*, World Resources Institute, Washington, D.C. (March).



UGH!
WISH I
FELT
BETTER!

Riley, B. & Cox, C. 1999, *Landscaping Nature's Way: Using Natural Landscaping to Reduce Herbicide Use*, NCAP, PO Box 1393 Eugene, OR 97440-1393 USA (www.pesticide.org).

Savage, E.P., Keefe, T.J. & Mounce, L.M. 1988, 'Chronic neurological sequelae of acute organophosphate poisoning', *Arch. Environ. Health*, 43:38-45.

Sawada, Y.Y., et al 1988, 'Probable toxicity of surface-active agent in commercial herbicide containing glyphosate', *Lancet*, 1(8580):299.

Solomon, G.M. & Mott, L.M. 1998, *Trouble On The Farm: Growing Up With Pesticides In Agricultural Communities*, Natural Resources Defense Council, New York, NY, USA.

Steenland, K; Jenkins, B; Ames, R.G.; O'Malley, M; Chrislip, D; Russo, J. 1994, 'Chronic neurological sequelae to organophosphate pesticide poisoning', *Am. Journal Public Health*, 84:731-736.

Stedman's Medical Dictionary 1976, 'Hypotonia', in *Stedman's Medical Dictionary*, 23rd Edn, The Williams & Wilkins Company, Baltimore Md USA.

Stephens, R; Sprugeon, A; Calvert, I.A.; Beach, J; Levy, L.S. Berry, H; Harrington, J.M. 1995, 'Neurophysiological effects of longterm exposure to organophosphates in sheep dip', *The Lancet* 345:1135-1139.

Talbot, A.R. 1991, 'Acute poisoning with a glyphosate-surfactant herbicide ('Roundup'): A review of 93 cases', *Human Exp. Toxicol.* 10:1-8).

Tandon, P; Padilla, S; Barone, S; Pope, C.N.; Tilson, H.A. 1994, 'Fenthion produces a persistent decrease in muscarinic receptor function in the adult rat retina', *Toxicol Appl Pharmacol*, 125:271-280.

Tominack, R.L. 1991, 'Taiwan national poison center survey of glyphosate-surfactant herbicide ingestions', *Clin. Toxicol.* 29(1):91-109.

Tortensson, N.T.L., Lundgren, L.N., and Stenstrom, J. 1989, 'Influence of climate and edaphic factors on persitence of glyphosate and 2,4-D in forest soils', *Ecotoxicol. Environ. Safety* 18:230-239.



Tortora, G.J., Funke, B.R. & Case, C.L. 1995, *Microbiology – An Introduction*, 5th edition, The Benjamin/Cummings Publishing Company, Inc., 390 Bridge Parkway.

US EPA Office of Pesticide Programs 1986, *Guidance for the reregistration of pesticide products containing glyphosate as the active ingredient*, Washington, D.C. (June).

US EPA 1990-1, *Nonoccupational pesticide exposure study (NOPES) Final Report*, EPA Office of Research and Development, Washington, DC (March).

US EPA 1993-2, 'Science chapter for registration eligibility document for glyphosate', EPA Ecological Effects Branch, Washington, DC (May 1).

US EPA 1994, 'Reregistration eligibility decision (RED): Oryzalin', Office of Prevention, Pesticides and Toxic Substances, Washington, DC (September).

US EPA 1999-1, Memorandum: 'Office of Pesticide Programs list of chemicals evaluated for carcinogenic potential', Office of Prevention, Pesticides and Toxic Substances, Washington, DC (August).

Vigfusson & Vyse 1980, 'The effect of the pesticides Dexon, Captan and Roundup on sister chromatid exchanges in human lymphocytes in vitro', *Mutation Research* 79:53-57.

Watanabe, et al, 1990, 'Placental and blood-brain barrier transfer following prenatal and postnatal exposures to neuroactive drugs: relationship with partition coefficient and behavioral teratogenesis', *Toxicol. Appl. Pharmacol.*, 105(1):66-77.

Williams & Eagle 1979, 'Persistence of dichlobenil in a sandy soil and effects of residues on plant growth', *Weed Res*, 19:315-319.

World Health Organization 1994, *Glyphosate*, Environmental Health Criteria #159, United Nations Environment Programme, Geneva, Switzerland.

Zweiner, R. & Ginsburg, C. 1988, 'Organophosphate and carbamate poisoning in infants and children', *Pediatrics* 81:121-126.